file: 53039-1_front_art
folder: 53039-1_TechPkg_Sevin 32oz
colors: 4cp, 186, white, rota-motion
☐ dp proofed_____

created by: customer
alt: 1-25-12 bm
size: 4.7566"(w) x 6.7122"(h)

Component #:	N/A - 201310
Description:	Sevin RTU 32oz - Front 32oz Label Front
UPC #:	N/A N/A
Printer:	CCL Label Memphis
Program:	Adobe Illustrator CS5
Creation Date:	10-03-13
Rev. Date:	12-19-13
Operator:	DMU

Ink Colors **Schawk Job No: 091712** **Item: A**

Yellow 1 2 25 50 75 100
PMS 186C 1 2 25 50 75 100
Cyan 1 2 25 50 75 100
Opaque White 1 2 25 50 75 100
Magenta 1 2 25 50 75 100
Black 1 2 25 50 75 100

SCHAWK!
2626 2nd Street NE,
Minneapolis, MN
55418 USA
612 789 8514
schawk.com

The color on this copy will not match the final printed package.
Schawk Minneapolis has checked this artwork for accuracy. Final approval is the client's responsibility.
Please double check for errors before reproduction.

UPC Check	Proofread/QC	QR Code
Initial Date	Proofreader Initial Date	Initial Date
Schawk Minneapolis		Schawk Minneapolis
Numerical Scan Only ☐ Pass ☐ Fail	Quality Control Initial Date	☐ Pass ☐ Fail

52926-2_base_art 2/28/12 1:55 PM Page 1

GARDENTECH
Sevin®

Ready-To-Spray Bug Killer

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product. Wear a hat and eye protection when making overhead applications. Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use. Wash the outside of gloves before removing. Wash thoroughly after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID
This product is a N-methyl carbamate that inhibits cholinesterase

IF SWALLOWED:	• Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 800-420-9347 for emergency treatment information.

NOTE TO PHYSICIAN: Carbaryl is a N-methyl carbamate that inhibits cholinesterase. Atropine is antidotal. Do not use 2-PAM (pyridene-2-aldoxime methiodide), opiates or cholinesterase inhibiting drugs. Probable mucosal damage may contraindicate the use of gastric lavage. To aid in the confirmation of diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analysis will be arranged by TechPac, LLC. Consultation on therapy can be obtained at all hours by calling the appropriate emergency number 800-420-9347 (U.S.A.)

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or allow drift to water. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. **BEE CAUTION: MAY KILL HONEYBEES AND OTHER BEES.** This product is highly toxic to honeybees and other bees. Do not apply to plants in bloom.

STORAGE AND DISPOSAL

STORAGE
Store unused GardenTech® Sevin® Ready To Spray Bug Killer in original container only, in cool, dry area out of reach of children and animals, preferably in a locked storage area. Do not store in areas where temperatures frequently exceed 100°F.
Disposal: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. **If partly filled:** Call your local solid waste agency or 866-945-5033 for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARNING: This product contains carbaryl, a chemical known to the State of California to cause cancer and reproductive harm in laboratory animals.

Distributed by: TechPac, LLC, 1000 Parkwood Circle, Ste. 700, Atlanta, GA 30339

EPA Reg. No. 264-334-71004
EPA Est. 67572-GA-1A, 67572-GA-2B
Subscript used is first letter of run code on container.

file: 52926-2_base_art
folder: 52926-2_Excel_Sevin 32oz
colors: black
☐ dp proofed_____

created by customer
alt: 2-28-12 bm
size: 4.0983"(w) x 6.7116"(h)

Component #: N/A - 201310
Description: Sevin RTU 32oz - Base 32oz Label Base
UPC #: N/A N/A
Printer: CCL Label Memphis
Program: Adobe Illustrator CS5
Creation Date: 10-03-13
Rev. Date: 1-07-14
Operator: DEL

Ink Colors Schawk Job No: 091712 Item: B
Black 1 2 25 50 75 100
Opaque White 1 2 25 50 75 100

SCHAWK!
2626 2nd Street NE, Minneapolis, MN 55418 USA
612 789 8514
schawk.com

The color on this copy will not match the final printed package. Schawk Minneapolis has checked this artwork for accuracy. Final approval is the client's responsibility. Please double check for errors before reproduction.

UPC Check: Initial Date; Schawk Minneapolis; Numerical Scan Only ☐ Pass ☐ Fail
Proofread/QC: Proofreader Initial Date; Quality Control Initial Date
QR Code: Initial Date; Schawk Minneapolis; ☐ Pass ☐ Fail

GARDEN TECH

Sevin®

Ready-To-Spray Bug Killer

FOR OUTDOOR RESIDENTIAL USE ONLY

For Vegetable Insects

Japanese Beetles

Horn/Cabbage Worms

Cutworms

Leafhoppers

Alfalfa Caterpillar, Armyworms, Aster Leafhopper, Bean Leaf Beetle, Colorado Potato Beetle, Cucumber Beetle, Cutworms, European Corn Borer, Fall Armyworm, Flea Beetles, Green Cloverworm, Harlequin Bug, Imported Cabbageworm, Japanese Beetle, Lace Bugs, Leafhoppers, Melon worm, Mexican Bean Beetle, Rednecked Peanutworm, Spittlebugs, Squash Bugs, Sweet Potato Weevil, Three Cornered Alfalfa Hopper, Thrips, Tomato Hornworm, Tomato Pinworm, Tortoise Beetles, Western Bean Cutworm, & many more.

Peel Back Tab for Additional Information

For Ornamental Pests

Moths

Grasshoppers

Tent Caterpillars

Japanese Beetles

Cicadas

Ticks*, Ants (excluding harvester, carpenter and pharaoh ants), Azalea Leafminer, Blister Beetle, Browntail Moth, Elm Leaf Aphid, Eriophyid Mites, Fall Armyworm, Fuller Rose Beetle, Plant Bugs, Scale Insects, Tent Caterpillars, & many more listed inside.

* Will kill pests that are present only at the time of applicatiion. Pest must be directly contacted for product to be effective.

EPA Reg. No. 264-334-71004
EPA Est. 67572-GA-1A, 67572-GA-2B
Subscript used is first letter of run code on container.

Product of China: Contains U.S. and Imported Ingredients

Distributed by:
TechPac, LLC
1000 Parkwood Circle
Suite 700
Atlanta, GA 30339

6 13499 07300 6
120210GGG

FIRST AID

This product is a N-methyl carbamate that inhibits cholinesterase

IF SWALLOWED:	• Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 800-420-9347 for emergency treatment information.

NOTE TO PHYSICIAN: Carbaryl is a N-methyl carbamate that inhibits cholinesterase. Atropine is antidotal. Do not use 2-PAM (pyridene-2-aldoxime methiodide), opiates or cholinesterase inhibiting drugs. Probable mucosal damage may contraindicate the use of gastric lavage. To aid in the confirmation of diagnosis, urine samples should be obtained within 24 hours of exposure and immediately frozen. Analysis will be arranged by TechPac, LLC. Consultation on therapy can be obtained at all hours by calling the appropriate emergency number 800-420-9347 (U.S.A.)

For MEDICAL Emergencies ONLY Call 24 Hours A Day 800-420-9347
For TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 800-424-9300
For PRODUCT USE Information Call 866-945-5033

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist. Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product. Wear a hat and eye protection when making overhead applications. Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use. Wash the outside of gloves before removing. Wash thoroughly after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or allow drift to water. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

BEE CAUTION: MAY KILL HONEYBEES AND OTHER BEES. This product is highly toxic to honeybees and other bees. Do not apply to plants in bloom.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

Strictly observe restrictions, precautions, and label directions.

RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- For outdoor use only.
- Not for use on plants grown for sale, commercial use, research, or for commercial seed production.
- Use on lawns is prohibited.

STORAGE AND DISPOSAL

STORAGE

Store unused GardenTech® Sevin® Ready To Spray Bug Killer in original container only, in cool, dry area out of reach of children and animals, preferably in a locked storage area. Do not store in areas where temperatures frequently exceed 100°F.

Disposal: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available.

If partly filled: Call your local solid waste agency or 866-945-5033 for disposal instructions. Never place unused product down any indoor or outdoor drain.

PLANT RESPONSE PRECAUTIONS

Application to wet foliage or during periods of high humidity may cause injury to tender foliage. Do not use on Boston Ivy, Virginia creeper and maidenhair fern as injury may result. Sevin® spray may also injure Virginia and sand pines.

STAINING PRECAUTION

Staining may occur on certain surfaces such as stucco, brick, cinder block, wood, fabrics, and carpet. Spray deposits that could inadvertently reach painted or stained surfaces or finishes (i.e., cars, houses, trailers, boats, etc.) should be immediately removed by washing to prevent discoloration. Avoid applications to surfaces where visible spray residues are objectionable. Tracking of product residues onto fabric or carpet may cause permanent discoloration.

SPECIFIC USE DIRECTIONS

HOSE-END SPRAYER INSTRUCTIONS

This ready to spray product delivers a spray solution containing approximately 0.45% carbaryl.

HOW TO USE	• Apply when insects or damage appear. Repeat as necessary to kill insects, following the specified spray interval listed in the specific directions below. • Apply when air is calm. • Shake bottle well before use. • Attach garden hose to spray nozzle. • Turn on water supply. • Push safety tab and turn control knob to "Water" position. • Point sprayer nozzle toward targeted plants and turn control knob to "On". Product will automatically mix with water. • Start spraying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. • Spray upper and lower leaf surfaces thoroughly. • Allow spray to dry in treated areas before reentering. • Do not apply to large trees. For trees taller than 10 feet, consider hiring a licensed professional.

GARDEN VEGETABLES

To effectively kill the vegetable pests listed below, apply product in adequate volume to effectively cover both the upper and lower surfaces of the plant.

PHI refers to the minimum interval in days that must be observed between the date of the last application and date of harvest. APP refers to the maximum number of applications per crop per year.

Vegetable	PHI	APP
Asparagus	1	3
Brassica head and stem vegetables*: Broccoli, Brussels sprouts, Cabbage, Cauliflower, Chinese cabbage, Kohlrabi	3	4

(continued)

GARDEN VEGETABLES *(continued)*

Vegetable	PHI	APP
Brassica leafy green vegetables*: Chinese cabbage, Collards, Kale, Mustard greens, Mustard spinach, Turnip greens	14	4
Corn (sweet)	2	8
Cucurbit vegetables: Cucumber, Melons, Pumpkin, Squash	3	6
Dried shelled legume vegetables**: Adzuki bean, Blackeyed pea, Cowpea, Field bean, Kidney bean, Lentils, Lima been (dry), Mung bean, Navy bean, Pinto bean, Southern pea	21	4
Edible-podded legume vegetables**: Dwarf pea, Edible-pod pea, Jack bean, Snap bean, Snow pea, Sugar snap pea, Wax bean, Yardlong bean.	3	4
Fruiting vegetables: Eggplant, Groundcherry, Okra, Peppers, Tomatoes	3	7
Leafy vegetables: Cardoon, Celery, Celtuce, Florence fennel, Dandelion, Endive, Lettuce (head and leaf), Parsley, Rhubarb, Spinach, Swiss chard	14	5
Root and Tuber vegetables (except sugar beets and sweet potatoes): Artichoke (Chinese and Jerusalem), Carrots, Garden Beets (roots and tops), Ginger, Ginseng, Horseradish, Parsnip, Potato, Radish, Rutabega, Salsify, Turnip (roots and tops), Yams	7	6
Sweet Potatoes	7	8
PESTS Alfalfa Caterpillar, Apache cicada, Armyworms, Asparagus beetle, Aster leafhopper, Bean leaf beetle, Chinch bug, Colorado potato beetle, Corn earworm, Corn rootworm Adult , Cowpea curculio, Cucumber beetle, Cutworms, European corn borer, Fall armyworm, Flea beetles, Grasshoppers, Green Cloverworm, Harlequin bug, Imported cabbageworm, Japanese beetle, Lace bugs, Leafhoppers, Lima Bean pod Borer, Lygus bugs, Melon worm, Mexican bean beetle, Pea weevil, Rednecked peanutworm, Sap Beetles, Southwestern corn borer, Spittlebugs, Squash bugs, Stink bugs, Sweet potato weevil, Three cornered alfalfa hopper, Thrips, Tomato hornworm, Tomato pinworm, Tortoise beetle, Velvetbean Webworms, Western bean Cutworm, Whitefringed Beetle (adults), Yellowstriped Armyworm		

* Application is only permitted within 30 days of crop emergence or the date of transplanting.

** Do not use on fresh/succulent peas and beans.

FRUITS

This product may be used as a spot treatment for localized pest infestations on the tree fruits listed below. Do not use on large trees or where pest populations are generally inaccessible to spray deposits, as inadequate performance will result.

Spray upper and lower leaf surfaces and between fruit clusters and on small limbs and trunks to the point of runoff where pest appear. Repeat applications as necessary up to a total of eight times per year for tree fruits or five times per year for small fruits, but not more often than once every seven days.

NOTE: To avoid undesired apple thinning delay use until at least 30 days after full bloom. Carbaryl insecticide may injure early dawn and sunrise strawberry varieties on the Delmarva Peninsula. The number in parenthesis following each fruit and nut crop grouping refers to the minimum interval that must be observed between the date of the last application and the date of harvest.

TREE FRUITS	PESTS
Apples, Apricots, Cherries, Nectarines, Peaches, Pears, Plums, Prunes (3 days until harvest)	Apple aphid, Apple maggot, Apple mealybug, Apple pandemic, Apple rust mite, Apple sucker, Avocado Leafroller, Bagworms, Black Cherry Aphid, Black scale, Leafrollers, Lecanium scales, Lesser appleworm, Lesser peachtree borer, Meadow spittlebug, Omnivorous leaftier, Strawberry fruitworm, Orange tortrix

TREE FRUITS	PESTS
Orange, Grapefruit, Lemon, Lime, Citrus Hybrids (e.g., Tangelo), Mandarin (Tangerine), Pummelo, Calamondin, Citron, Kumquat (5 days until harvest)	Apopka weevil (adult), Avocado leafroller, California orangedog, Citrus cutworm, Citrus root weevils (adults), Citrus rust mite, Eriophyid mites, Fruittree leafroller, Fuller rose beetle, Little leaf notcher (adult), Orange tortrix, Plant bugs, Scale insects [Black scale, brown soft scale, California red scale (except in California), citrus snow scale, yellow scale (except in California)], Western tussock moth
Small Fruits Blackberries, Blueberries, Dewberries, Grapes, Raspberries, Strawberries (7 days Until harvest)	Blueberry maggot, Brown soft scale, California red scale, Cherry fruitworm, Cherry maggot, Citrus cutworm, Citrus root weevil, Citricola scale, Citrus snow scale, Codling moth, Cucumber beetles, Cutworms, Eastern tent caterpillar, Eightspotted forester, European apple sawfly, European earwig, European raspberry aphid, Eyespotted budmoth, Forbes scale, Fruitree leafroller, Grape berry moth, Grape leaffolder, Grape leafhopper, Green fruitworm, Japanese beetle, June beetles, Leafhoppers, Oriental fruit moth, Oystershell scale, Peach twig borer, Pear leaf blister mite, Pear psylia, Pear rust mite, Periodical cicada, Plum curculio, Prune leafhopper, Raspberry sawfly, Red-banded leafroller, Rose chafer, Rosy apple aphid, Saltmarsh caterpillar, San Jose scale, Snowy tree cricket, Fireworm, Strawberry weevil, Tarnished plant bug, Tentiform leafminers, Tussock moth, Variegated leafroller, Western grapeleaf skeletonizer, Western tussock moth, West Indian sugarcaneborer (adults), Western yellowstripe armyworm, White apple leafhopper, Wooly apple aphid, Yellow headed fireworm

ORNAMENTAL TREES, SHRUBS AND FLOWERS

This product may be used as a spot treatment for localized pest infestations on trees. Do not use on large trees or where pest populations are generally inaccessible to spray deposits.

Spray upper and lower leaf surfaces and small trunks, stems and twigs to the point of runoff where pests appear. Repeat applications as necessary up to a total of four times per year for trees and six times per year for ornamental shrubs and flowers, but not more often than once every seven days.

ORNAMENTAL TREES, SHRUBS AND FLOWERS	PESTS
Ornamental Trees, Shrubs and Flowers	Ants (excluding harvester, carpenter, and pharoah ants), Apple aphid, Armyworm, Azalea leafminer, Bagworms, Birch leafminer, Blister beetle, Boxelder bug, Boxwood leafminer, Browntail moth, Cankerworms, Catalpa sphinx, Chiggers*, Cooley spruce gall, Elm leaf aphid, Elm spanworm, Eriophyid mites, European pine-shoot moth, Fall armyworm, Flea beetles, Fuller rose beetles, Gall midges, Gall wasps, Grasshoppers, Green striped-Mapleworm, Gypsy moth, Hackberry nipplegall makers, Holly budmoth, Holly leafminer, Jackpine budworm, Japanese beetle, Jeffrey pine needleminer, June beetles, Plant bugs, Lace bugs, Leafhoppers, Leafrollers, Locust borer, Maple leafcutter, Mealy bugs, Mimosa webworm, Nantucket pine tip moth, Oak leafminers, Oak leaf skeletonizer, Oakworm complex, Oleander caterpillar, Olive ash borer, Orange-striped oakworm, Orange tortrix, Periodical cicada, Pine Sawfly, Pine spittlebug, Pitch pine tip moth, Poinsettia horn worm, Psyllids, Puss caterpillar, Redhumped oakworm, Rose aphid, Rose chafer, Roseslug, Saddled prominent, Sawflies (exposed), Scale insects, Sowbugs, Spiny elm caterpillar, Springtails, Spruce budworm, Spruce needleminer, Subtropical pine tip moth, Tent caterpillar, Thorn bug, Thrips (exposed), Ticks*, Walnut caterpillar-Webworms, Western hemlock loopers, Western spruce budworm, Willow leaf beetles, Yellow poplar-weevil

* Will kill pests that are present only at the time of application. Pest must be directly contacted for product to be effective.

NUISANCE PESTS AROUND OUTDOOR RESIDENTIAL AREAS

This product may be used as a perimeter treatment to kill listed nuisance pests. Thoroughly wet the outside perimeter of the home where pests tend to congregate. Use on lawns is prohibited. Will kill target pests that are present only at the time of application. Pest must be directly contacted for product to be effective.

NOTE: Staining may occur on certain surfaces such as stucco, brick, cinder block and wood. Therefore, applications of this product to surfaces where a noticeable residue or discoloration is objectionable should be avoided.

PESTS
Ants (except pharaoh, harvester, and carpenter ants), Bees, Centipedes, Cockroaches, Crickets, Darkling beetles, Earwigs, Firebrats, Fleas, Grasshoppers, Millipedes, Mosquitoes (adults), Scorpions, Silverfish, Spiders, Ticks, and Wasps

NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Sevin is a registered trademark of Tessenderlo Kerley, Inc.

GardenTech is a registered trademark of Gulfstream Home & Garden, Inc.

PRIMEROS AUXILIOS

Este producto es un N-metil carbamato que inhibe la colinesterasa.

EN CASO DE SER INGERIDO:	• Llame inmediatamente a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento. • No induzca el vómito, a menos que se lo indique un médico o el centro de control de envenenamientos. • Haga que la persona tome agua, si puede tragar. • No administre nada por boca a una persona que haya perdido el conocimiento.
SI ESTÁ EN LA PIEL O EN LA ROPA:	• Quítese inmediatamente la ropa contaminada. • Enjuague la piel inmediatamente con abundante agua entre 15 a 20 minutos. • Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.
SI SE INHALA:	• Lleve la persona a un lugar con aire fresco. • Si la persona no respira, llame al 911 o a una ambulancia, luego administre respiración artificial, preferentemente boca a boca, si es posible. • Llame a un centro de control de envenenamientos o a un médico para recibir más instrucciones de tratamiento.
SI ESTÁ EN LOS OJOS:	• Mantenga el ojo abierto y enjuague lenta y suavemente con agua durante 15 a 20 minutos. • Si lleva lentes de contacto puestos, retírelos luego de los primeros 5 minutos, luego continúe enjuagando. • Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.

Cuando llame a un centro de control de envenenamientos o a un médico, o busque tratamiento, tenga a mano el envase o la etiqueta del producto. También puede ponerse en contacto con el 800-420-9347 para obtener información de tratamiento de emergencia

NOTA PARA EL MÉDICO: El Carbarilo es un N-metil carbamato que inhibe la colinesterasa. La atropina es un antídoto. No use 2-PAM (piridina-2-aldoxima metiodida), opiáceos o medicamentos inhibidores de la colinesterasa. posible daño de la mucosa, lo que podría contraindicar un lavado gástrico. Para ayudar en la confirmación del diagnóstico, se deben obtener muestras de orina dentro de 24 horas de la exposición y congelarlas inmediatamente. TechPac, LLC. coordinará el análisis. Se pueden realizar consultas sobre terapia las 24 horas llamando al número de emergencias correspondiente 800-420-9347 (EE.UU.)

SÓLO para emergencias médicas llame las 24 horas al 800-420-9347

SÓLO para emergencias durante el transporte llame las 24 horas al 800-424-9300

Para obtener información sobre USO DEL PRODUCTO llame al 866-945-5033

AVISOS DE PRECAUCIÓN

RIESGOS PARA HUMANOS Y ANIMALES DOMÉSTICOS

PRECAUCIÓN

El producto es nocivo si se ingiere. Es dañino si se absorbe a través de la piel. El producto es nocivo si se inhala. Evite el contacto con la piel, los ojos o la ropa. Evite respirar los vapores o el rocío. Al mezclar y aplicar este producto, use camisa de mangas largas, pantalones largos, medias y zapatos y guantes de látex o goma. Use casco y lentes de protección cuando lo aplique en altura. Quítese la ropa inmediatamente si ésta se empapa del pesticida. Cámbiese la ropa lo antes posible una vez terminada su aplicación. Lave el exterior de los guantes antes de retirarlos. Lávese bien después de manipular el producto y antes de comer, beber, masticar chicle, consumir tabaco o usar el inodoro.

ADVERTENCIA: Este producto contiene carbarilo, una sustancia química reconocida por el estado de California como causante de cáncer y daños en el aparato reproductivo en animales de laboratorio.

RIESGOS PARA EL MEDIO AMBIENTE
Este producto es extremadamente tóxico para los invertebrados acuáticos. No aplique directamente en el agua, ni permita que se escurra hasta el agua. Para proteger el medio ambiente, no deje que el pesticida entre o escurra hacia desagües pluviales, zanjas de desagüe, alcantarillas ni aguas superficiales. Aplicar este producto en un clima tranquilo, cuando no haya pronóstico de lluvia durante las siguientes 24 horas, ayudará a asegurar que el viento o la lluvia no soplen o laven el pesticida del área de tratamiento. Enjuagar el equipo de aplicación sobre el área tratada ayudará a evitar los escurrimientos hacia masas de agua o sistemas de desagüe.
PRECAUCIÓN CON LAS ABEJAS: PUEDE MATAR ABEJAS PRODUCTORAS DE MIEL Y OTROS TIPOS DE ABEJAS. Este producto es extremadamente tóxico para las abejas productoras de miel y otros tipos de abejas. No aplique a las plantas en floración.

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto con fines que no sean los indicados en la etiqueta.
Lea la etiqueta completamente antes de utilizar este producto.
Cumpla estrictamente con las restricciones, precauciones e instrucciones de la etiqueta.

RESTRICCIONES

- No aplique el producto de ninguna forma que pueda entrar en contacto con personas o mascotas, ya sea directa o indirectamente. Mantenga a las personas y mascotas alejadas del área durante la aplicación.
- No permita que las personas o mascotas ingresen al área tratada hasta que el aerosol se haya secado.
- Sólo para uso en exteriores.
- No lo use en plantas cultivadas para la venta, uso comercial, para fines de investigación o la producción comercial de semillas.
- Se prohíbe su uso en céspedes.

ALMACENAJE Y ELIMINACIÓN

ALMACENAJE
Almacene el insecticida GardenTech Sevin listo para usar sin usar sólo en su envase original en un lugar fresco, seco y fuera del alcance de los niños y animales, preferiblemente en un área de almacenaje cerrada con llave. No lo almacene en áreas donde la temperatura supere frecuentemente los 37,8 °C.
Eliminación: Si está vacío: El envase no se puede rellenar. No vuelva a usar ni rellene este envase. Ofrézcalo para reciclaje si está disponible.
Si está parcialmente lleno: Llame a su agencia local de eliminación de desechos sólidos o al 866-945-5033 para obtener instrucciones sobre cómo desechar el producto. Nunca deseche el producto sin usar en desagües en interiores o exteriores.

PRECAUCIONES DE USO EN PLANTAS
La aplicación en follaje húmedo o durante períodos de alta humedad puede causar daños al follaje tierno. No utilizar en Hiedra de Boston, Parra de Virginia, ni en culantrillo ya que podría dañarlos. Sevin también puede dañar los pinos de las arenas de Virginia.

PRECAUCIONES POR MANCHAS
Ciertas superficies se pueden manchar, tales como las de estuco, ladrillo, bloques de hormigón, madera, tela y alfombra. Los depósitos del rociado que pueden alcanzar inadvertidamente superficies o acabados pintados o teñidos (es decir, automóviles, casas, remolques, botes, etc.) deben ser retirados inmediatamente por medio de lavado para evitar la decoloración. Evite la aplicación en superficies en las que los residuos de rociado visibles no sean tolerables. Las marcas de residuos del producto sobre telas o alfombras pueden causar decoloración permanente.

INSTRUCCIONES DE USO ESPECÍFICO

INSTRUCCIONES ATOMIZADOR PARA EXTREMO DE MANGUERA

Este producto listo para rociar proporciona una solución de rociado que contiene aproximadamente 0,45% de carbarilo.

CÓMO SE UTILIZA	• Aplique cuando aparezcan insectos o daños. Repita según sea necesario eliminar insectos, siguiendo el intervalo de rocío especificado en las instrucciones específicas a continuación. • Aplique cuando el aire esté calmo. • Agite bien la botella antes usar. • Conecte la manguera para jardín a la boquilla de rociado. • Abra el suministro de agua. • Presione la lengüeta de seguridad y gire la perilla a la posición "Water" (Agua). • Dirija la boquilla del atomizador hacia las plantas deseadas y gire la perilla de control a "On" (Encender). El producto se mezcla automáticamente con agua. • Comience la aplicación en la esquina más alejada del área a tratar y trabaje de atrás hacia delante para evitar el contacto con superficies húmedas. • Rocíe completamente las superficies superior e inferior de las hojas. • Deje secar el rocío en las áreas tratadas antes de volver a ingresar a ellas. • No aplique en árboles grandes. Para árboles cuya altura supere los 3,05 metros, considere la posibilidad de contratar un profesional autorizado.

VERDURAS DE HUERTO

Para eliminar eficientemente las plagas que atacan las verduras que se enumeran a continuación, aplique este producto en un volumen adecuado que permita cubrir tanto la superficie superior como la inferior de la planta. La sigla PHI se refiere al intervalo mínimo de días que debe respetar entre la fecha de la última aplicación y la cosecha. La sigla APP se refiere al número máximo de aplicaciones por cosecha por año.

Verdura	PHI	APP
Espárrago	1	3
Verduras de especie Brassica con cabeza y tallo*: Brócoli, repollitos de Bruselas, repollo, coliflor, repollo chino, colinabo	3	4
Verduras de especie Brassica de hojas verdes*: Repollo chino, col rizada, berzas, mostaza parda, espinaca mostaza, hojas de nabo	14	4
Maíz (dulce)	2	8
Verduras de la familia de las cucurbitáceas: Pepino, melones, zapallo, calabaza	3	6
Legumbres de vainas seca**: Frijol Adzuki, alubia, caupí, haba, frijol común, lentejas, judía de Lima (seca), frijol mungo, frijol Navy, frijol Pinto, arveja del sur	21	4
Legumbres de vainas comestibles**: Arveja enana, arveja de vaina comestible, canavalia, ejote, arveja china, tirabeque, frijol amarillo, frijol espárrago.	3	4
Verduras de frutos: Berenjena, camambú, quingombó, pimientos, tomates	3	7
Verduras de hoja: Cardo, apio, lechuga apio, hinojo de Florencia, diente de león, endivia, lechuga (cabeza y hojas), perejil, ruibarbo, espinaca, acelga	14	5
Verduras de raíz y tubérculos (excepto remolacha y camote): Alcachofa (china y de Jerusalén), zanahorias, remolacha común (raíces y cabezas), jengibre, ginseng, rábano picante, chirivía, papa, rábano, nabicol, salsifi, nabo (raíces y cabezas), boniato	7	6
Camotes	7	8

(continuación)

VERDURAS DE HUERTO *(continuación)*

PLAGAS
Oruga de la alfalfa, cigarra apache, gusano militar, escarabajo del espárrago, cigarra del áster, escarabajo de la hoja del frijol, chinche, escarabajo de la papa de Colorado, gusano del oído del maíz, crisomela del maíz adulta, curculio del caupí, escarabajo rayado del pepino, gusanos cortadores, piral del maíz, gusano militar de otoño, escarabajo pulga, saltamontes, gusanos verdes del trébol, chinche arlequín, oruga de la col, escarabajo japonés, chinche de encaje, cigarras, gusano de las vainas, chinches lygus, gusano del melón, escarabajo mexicano del frijol, bruco de la arveja, gusano cuello rojo del maní, escarabajos de la fruta seca, barrenador del maíz del Suroeste, cigarra espumadora, chinches de la calabaza, chinches verdes de campo, tetuán del boniato, periquito tricornudo, trips, gusano de cuerno del tomate, polilla del tomate, escarabajo tortuga, oruga del nogal, gusano de bolsa, gusano cortador occidentales del frijol, escarabajo franjado blanco (adultos), gusano militar de rayas amarillas

*Sólo se permite su aplicación dentro de los 30 días desde que los cultivos emergen o de la fecha del trasplante.

**No use en arvejas y frijoles frescos/suculentos.

FRUTAS

Este producto se puede usar como tratamiento localizado para infestaciones por plagas ubicadas en las frutas de los árboles mencionados a continuación. No use en árboles grandes o en aquellos lugares donde comúnmente las plagas son inaccesibles para los aerosoles, puesto que no se logrará un desempeño adecuado.

Rocíe hasta que escurra directamente en las superficies superiores e inferiores de las hojas, entre los racimos de frutas y en las ramas y troncos pequeños donde aparezcan plagas. Repita las aplicaciones según sea necesario hasta un total de ocho veces por año para las frutas de árboles, o cinco veces por año para las frutas pequeñas, pero no más seguido que una vez cada siete días.

NOTA: Para evitar el debilitamiento no deseado de las manzanas, no lo utilice hasta al menos 30 días después de la floración plena. El carbarilo es un insecticida que puede dañar las variedades de fresas Early Dawn y Sunrise en la Península de Delmarva. La cifra indicada entre paréntesis después de cada grupo de cultivo de frutas y frutos secos se refiere al intervalo mínimo que se debe respetar entre la fecha de la última aplicación y la fecha de la cosecha.

FRUTAS DE ÁRBOLES	PLAGAS
Manzanas, albaricoques, cerezas, nectarinas, melocotones, peras, ciruelas, ciruelas pasas (3 días hasta la cosecha)	Áfido del manzano, gusano del manzano, gorgojos del manzano, pandemis del manzano, ácaros del polvo del manzano, áfido azul del manzano, orugas enrolladoras del aguacate, gusanos de saco, áfido negro del cerezo, escama negra, orugas enrolladoras de las hojas, escamas Lecanium, gusano menor del manzano, barrenador menor del melocotón, salivazos de los pastos, tórtrix omnívora de las hojas, gusano de las fresas, tórtrix de la naranja
Naranja, pomelo, limón, híbridos cítricos (por ejemplo, tangelo), mandarina, calamondín, cidra, quinoto (5 días hasta la cosecha)	Escarabajo de Apopka (adulto), oruga enrolladora del aguacate, oruga de la mariposa de California, oruga cortadora de cítricos, escarabajo rubio de la raíz de cítricos (adulto), ácaros del polvo de los cítricos, ácaros eriofidos, oruga enrolladora de árboles frutales, escarabajo de la rosa Fuller, pequeño comedor de hojas (adulto), tórtrix de la naranja, insectos de plantas, insectos escama [escama negra, escama café, escama roja de California (excepto en California), escama cítrica de nieve, escama amarilla (excepto en California)], polilla occidental de manchas blancas

(continuación)

FRUTAS PEQUEÑAS	PLAGAS
Moras, arándanos, zarzas, uvas, frambuesas, fresas (7 días hasta la cosecha)	Gusano del arándano, escama café, escama roja de California, gusano de la fruta del cerezo, gusano del cerezo, oruga cortadora de cítricos, escarabajo rubio de la raíz de cítricos, escama citrícola, oruga cortadora de cítricos, escama cítrica de nieve, agusanado de la manzana, escarabajo del pepino, orugas cortadoras, oruga tendera del este, mariposa nocturna de ocho manchas, mosca de sierra europea de la manzana, tijeretas europeas, áfido europeo de la frambuesa, polilla roja de las yemas, escama de Forbes, orugas enrolladoras de las hojas de los frutales, polilla de la uva, oruga enrolladora de la hoja de la uva, cigarra de la uva, gusano verde de la fruta, escarabajo japonés, escarabajo de junio, cigarra, polilla oriental de la fruta, escama del carapacho de ostra, barrenador de ramas del melocotón, ácaro de ampolla de la hoja del peral, psila del peral, ácaro del polvo del peral, cigarra periódica, curculio de la ciruela, cigarra de las pasas, mosca de sierra de la frambuesa, oruga enrolladora con banda roja, abejorro de la rosa, áfido de la manzana rosácea, oruga peluda, escama de San José, grillo de Fulton, gusanos de elatéridos, escarabajo rubio de la frambuesa, insecto descolorado de plantas, minadores de hojas, polilla de manchas blancas Tussock, oruga enrolladora de hojas variegadas, oruga occidental de la parra, polilla occidental de manchas blancas, barrenador de caña de azúcar india (adultos), gusano soldado occidental de rayas amarillas, cigarra blanca del manzano, áfido del manzano velloso, elatérido de cabeza amarilla

ÁRBOLES ORNAMENTALES, ARBUSTOS Y FLORES

Este producto se puede usar como tratamiento localizado para infestaciones por plagas ubicadas en los árboles. No use en árboles grandes o en aquellos lugares donde comúnmente las plagas son inaccesibles para los aerosoles.

Rocíe hasta que escurra directamente en las superficies superiores e inferiores de las hojas y en las ramas, tallos y troncos pequeños donde aparezcan plagas. Repita las aplicaciones según sea necesario hasta un total de cuatro veces por año para los árboles o seis veces por año para los arbustos y flores ornamentales, pero no más seguido que una vez cada siete días.

ÁRBOLES ORNAMENTALES, ARBUSTOS Y FLORES	PLAGAS
Árboles ornamentales, arbustos y flores	Hormigas (excluyendo las hormigas cosecheras, carpinteras y faraón), áfido de la manzana, gusano soldado, minadores de hojas de la Azalea, gusanos del saco, minadores de hojas del abedul, escarabajo de ampolla, chinche del arce americano, minadores de hojas del Boj, polilla cola café, gusanos medidores, ceratomia catalpae, niguas*, adelges Cooley, áfido de la hoja del olmo, ennomo del olmo, ácaros eriofidos, polilla europea del brote del pino, gusano soldado de otoño, pulguillas, escarabajos de la rosa Fuller, mosquitos pequeños, avispas pequeñas, saltamontes, gusano del arce de rayas verdes, polilla gitana, pulguillas del álmez, polilla roja del acebo, minador de hojas del acebo, gusanos del pino de Jack, escarabajo japonés, minador de brotes del pino Jeffrey, escarabajo de junio, insectos de plantas, escamas, cigarras, orugas enrolladoras de hojas, barrenador de algarrobo, cortador de hojas del arce, gorgojos, orugas tejedoras de la mimosa, polilla de Nantucket de brotes del pino, minadores de hojas del roble, oruga de la hoja del roble, gusano complejo del roble, oruga de la adelfa, barrenador de fresno, gusano de roble con rayas anaranjadas, tórtrix de la naranja, cigarra periódica, mosca de sierra del pino, salivazo del pino, polilla de brote de pino rígido, gusano con cuernos de la poinsettia, pulguillas,

ÁRBOLES ORNAMENTALES, ARBUSTOS Y FLORES *(continuación)*

ÁRBOLES ORNAMENTALES, ARBUSTOS Y FLORES	PLAGAS
Árboles ornamentales, arbustos y flores	oruga peluche, gusano rojo del roble, áfido de la rosa, abejorro de la rosa, babosa de la rosa, oruga ensillada, moscas de cierra (expuestas), insectos escama, cochinillas, oruga espinosa del olmo, colémbolos, gusanos del abeto, minador de brotes del abeto, polilla subtropical de brote del pino, oruga tendera, insectos espina, pulgones (expuestos), garrapatas*, orugas nogales – orugas tejedoras, gusano medidor occidental del abeto Hemlock, gusanos del abeto occidental, escarabajo de las hojas del sauce, escarabajo amarillo del álamo

Sólo eliminará las plagas presentes al momento de la aplicación. Para que el producto sea efectivo, se debe aplicar directamente sobre la plaga.

PARA USAR EN PLAGAS MOLESTAS PRESENTES EN ÁREAS RESIDENCIALES EN EXTERIORES

Este producto se puede usar como tratamiento perimetral para eliminar las plagas molestas mencionadas. Empape cuidadosamente el perímetro exterior del hogar donde se tienden a acumular las plagas. Se prohíbe su uso en céspedes. Eliminará las plagas específicas que se encuentren presentes al momento de la aplicación. Para que el producto sea efectivo, se debe aplicar directamente sobre la plaga.

NOTA: Ciertas superficies se pueden manchar, tales como las de estuco, ladrillo, bloques de hormigón y madera. Por lo tanto, se debe evitar aplicar este producto sobre las superficies donde no se desea observar residuos o decoloración notorios.

PLAGAS
Hormigas (excepto la hormiga faraón, cosechera y carpintera), escarabajos, ciempiés, cucarachas, grillos, escarabajos negros, tijeretas, termobias, pulgas, saltamontes, milpiés, mosquitos (adultos), escorpiones, lepismas, arañas, garrapatas y avispas.

AVISO: Los estudios y las pruebas han determinado que el "modo de empleo" es adecuado para el uso correcto de este producto bajo las condiciones esperadas. En la medida que lo permita la ley vigente, el comprador asume la responsabilidad ante un mal desempeño o la falta de seguridad si no se respetan las instrucciones de uso.

Sevin es una marca comercial registrada de Tessenderlo Kerley, Inc.
GardenTech es una marca comercial registrada de Gulfstream Home & Garden, Inc.